

## CMA 120 System for Freely Moving Animals

1. Paasonen, J. et al., 2017. Dose-response effect of acute phencyclidine on functional connectivity and dopamine levels, and their association with schizophrenia-like symptom classes in rat. *Neuropharmacology*, 119, pp.15–25.
2. Jin, W.-S. et al., 2017. Peritoneal dialysis reduces amyloid-beta plasma levels in humans and attenuates Alzheimer-associated phenotypes in an APP/PS1 mouse model. *Acta Neuropathologica*, pp.1–14.
3. Bank, J.H.H. et al., 2017. Gene expression analysis and microdialysis suggest hypothalamic triiodothyronine (T3) gates daily torpor in Djungarian hamsters (*Phodopus sungorus*). *Journal of Comparative Physiology B*, pp.1–12.
4. Xiao, B.-X. et al., 2016. Pharmacokinetic profiles of the five isoflavonoids from *Pueraria lobata* roots in the CSF and plasma of rats. *Journal of Ethnopharmacology*, 184, pp.22–29.
5. Lindqvist, A. et al., 2016. In vivo Functional Evaluation of Increased Brain Delivery of the Opioid Peptide DAMGO by Glutathione-PEGylated Liposomes. *Pharmaceutical Research*, 33(1), pp.177–185.
6. He, Y. et al., 2016. In situ derivatization-ultrasound-assisted dispersive liquid–liquid microextraction for the determination of neurotransmitters in Parkinson’s rat brain microdialysates by ultra high performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 1458, pp.70–81.
7. Ortega, J., Meana, J.J. & Callado, L., 2016. In Vivo Brain Microdialysis of Monoamines. In R. Luján & F. Ciruela, eds. *Receptor and Ion Channel Detection in the Brain*. Neuromethods. Springer New York, pp. 415–434. Available at: [http://dx.doi.org/10.1007/978-1-4939-3064-7\\_25](http://dx.doi.org/10.1007/978-1-4939-3064-7_25) .
8. Qi, D. et al., 2016. Quantification of Dopamine in Brain Microdialysates with High-Performance Liquid Chromatography–Tandem Mass Spectrometry. *Analytical Sciences*, 32(4), pp.419–424.
9. Wang, F. et al., 2015. Sensitive Determination of Amantadine in Microdialysis Samples from Rat Plasma by HPLC with Fluorescence Detection. *Journal of Liquid Chromatography & Related Technologies*, 38(17), pp.1622–1628.
10. Keeler, G.D., Durdik, J.M. & Stenken, J.A., 2015a. Effects of delayed delivery of dexamethasone-21-phosphate via subcutaneous microdialysis implants on macrophage activation in rats. *Acta Biomaterialia*, 23, pp.27–37.
11. Wei, J. et al., 2015. Alteration of glutamate/GABA balance during acute alcohol intoxication in rats: Effect of Xingnaojing injection. *Journal of Ethnopharmacology*, 166, pp.333–339.
12. Alonso-Alonso, S. et al., 2015. Partial seizures induced by latrunculin A microperfusion in the mouse hippocampus: Role of extracellular glutamate and NMDA receptors. *Molecular & Cellular Epilepsy*, 2(0). Available at: <http://smartscitech.com/index.php/MCE/article/view/608> .
13. Keeler, G.D., Durdik, J.M. & Stenken, J.A., 2015b. Localized delivery of dexamethasone-21-phosphate via microdialysis implants in rat induces M(GC) macrophage polarization and alters CCL2 concentrations. *Acta Biomaterialia*, 12, pp.11–20.
14. Wang, Q. et al., 2014. Application of online microdialysis coupled with liquid chromatography-tandem mass spectrometry method in assessing neuroprotective effect of *Rhizoma coptidis* on diabetic rats. *Analytical Methods*, 7(1), pp.45–52.

15. Freire-Cobo, C. et al., 2014. The Calcineurin Inhibitor Ascomycin Interferes with the Early Stage of the Epileptogenic Process Induced by Latrunculin A Microperfusion in Rat Hippocampus. *Journal of Neuroimmune Pharmacology*, 9(5), pp.654–667.
16. Lietsche, J. et al., 2014. Self-built microdialysis probes with improved recoveries of ATP and neuropeptides. *Journal of Neuroscience Methods*, 237, pp.1–8.
17. Huo, T. et al., 2014. Effect of realgar on extracellular amino acid neurotransmitters in hippocampal CA1 region determined by online microdialysis–dansyl chloride derivatization–high-performance liquid chromatography and fluorescence detection. *Biomedical Chromatography*, 28(9), pp.1254–1262.
18. Netzahualcoyotzi, C. & Tapia, R., 2014. Energy Substrates Protect Hippocampus Against Endogenous Glutamate-Mediated Neurodegeneration in Awake Rats. *Neurochemical Research*, 39(7), pp.1346–1354.
19. Keeler, G.D., Durdik, J.M. & Stenken, J.A., 2014. Comparison of microdialysis sampling perfusion fluid components on the foreign body reaction in rat subcutaneous tissue. *European Journal of Pharmaceutical Sciences*, 57, pp.60–67.
20. Wang, Y., Chen, X., et al., 2014. Additional noradrenergic depletion aggravates forelimb akinesia and abnormal subthalamic nucleus activity in a rat model of Parkinson’s disease. *Life Sciences*, 119(1–2), pp.18–27.
21. Chen, X. et al., 2014. Effect of transporter inhibition on the distribution of cefadroxil in rat brain. *Fluids and Barriers of the CNS*, 11, p.25.
22. Sierra-Paredes, G. et al., 2014. Effects of eslicarbazepine acetate on acute and chronic latrunculin A-induced seizures and extracellular amino acid levels in the mouse hippocampus. *BMC Neuroscience*, 15, p.134.
23. Wang, Y., Wang, H.S., et al., 2014. L-DOPA-induced dyskinesia in a rat model of Parkinson’s disease is associated with the fluctuational release of norepinephrine in the sensorimotor striatum. *Journal of Neuroscience Research*, 92(12), pp.1733–1745.
24. Bonhomme, D. et al., 2014. Retinoic acid modulates intrahippocampal levels of corticosterone in middle-aged mice: consequences on hippocampal plasticity and contextual memory. *Frontiers in Aging Neuroscience*, 6. Available at: <http://journal.frontiersin.org/article/10.3389/fnagi.2014.00006/full> .
25. Alfaro-Rodriguez, A. et al., 2013. Histamine H3 receptor activation prevents dopamine D1 receptor-mediated inhibition of dopamine release in the rat striatum: A microdialysis study. *Neuroscience Letters*, 552, pp.5–9.
26. Fernández-Pastor, B., Ortega, J.E. & Meana, J.J., 2013. Involvement of serotonin 5-HT3 receptors in the modulation of noradrenergic transmission by serotonin reuptake inhibitors: a microdialysis study in rat brain. *Psychopharmacology*, 229(2), pp.331–344.
27. Tanda, G. et al., 2013. Relations between stimulation of mesolimbic dopamine and place conditioning in rats produced by cocaine or drugs that are tolerant to dopamine transporter conformational change. *Psychopharmacology*, 229(2), pp.307–321.
28. Grossman, R. et al., 2013. Microdialysis measurement of intratumoral temozolomide concentration after cediranib, a pan-VEGF receptor tyrosine kinase inhibitor, in a U87 glioma model. *Cancer Chemotherapy and Pharmacology*, 72(1), pp.93–100.
29. Ortega, J.E. et al., 2013. Additive effect of rimonabant and citalopram on extracellular serotonin levels monitored with in vivo microdialysis in rat brain. *European Journal of Pharmacology*, 709(1–3), pp.13–19.
30. Polissidis, A. et al., 2013. The cannabinoid CB1 receptor biphasically modulates motor activity and regulates dopamine and glutamate release region dependently. *International Journal of Neuropsychopharmacology*, 16(2), pp.393–403.
31. Rushaidhi, M. et al., 2013. Participation of hippocampal agmatine in spatial learning: An in vivo microdialysis study. *Neuropharmacology*, 65, pp.200–205.

32. Lin, L. et al., 2013. In vivo study on the monoamine neurotransmitters and their metabolites change in the striatum of Parkinsonian rats by liquid chromatography with an acetylene black nanoparticles modified electrode. *Journal of Pharmaceutical and Biomedical Analysis*, 72, pp.74–79.
33. Zielke, H.R., Zielke, C. & Baab, P., 2013. In Vivo Microdialysis for Measurement of Oxidative Metabolism in the Living Brain. In G. Di Giovanni et al., eds. *Microdialysis Techniques in Neuroscience*. Neuromethods. Humana Press, pp. 89–102. Available at: [http://dx.doi.org/10.1007/978-1-62703-173-8\\_5](http://dx.doi.org/10.1007/978-1-62703-173-8_5) .
34. Di Giovanni, G. et al., 2013. In Vivo Microdialysis to Study Striatal Dopaminergic Neurodegeneration. In G. Di Giovanni et al., eds. *Microdialysis Techniques in Neuroscience*. Neuromethods. Humana Press, pp. 23–42. Available at: [http://dx.doi.org/10.1007/978-1-62703-173-8\\_2](http://dx.doi.org/10.1007/978-1-62703-173-8_2) .
35. Ortega, J.E. et al., 2012. Regulation of central noradrenergic activity by 5-HT3 receptors located in the locus coeruleus of the rat. *Neuropharmacology*, 62(8), pp.2472–2479.
36. Sehba, F. & Bederson, J., 2012. Assessment of Global Ischemia. In J. Chen et al., eds. *Animal Models of Acute Neurological Injuries II*. Springer Protocols Handbooks. Humana Press, pp. 373–385. Available at: [http://dx.doi.org/10.1007/978-1-61779-576-3\\_25](http://dx.doi.org/10.1007/978-1-61779-576-3_25).
37. Chauveau, F. et al., 2010. Rapid stress-induced corticosterone rise in the hippocampus reverses serial memory retrieval pattern. *Hippocampus*, 20(1), pp.196–207.